

RNA BINDING PROTEIN AND BINDING SITEUSEFUL FOR EXPRESSION OF RECOMBINANT MOLECULES

5 *Inv* This is a national stage application filed under 35 USC 371, of PCT/US98/00840, filed 1/16/98. This Appn claims benefit of provisional 60/035,955 01/07/97 Technical Field and provisional appn 60/069,400 12/12/97

The invention relates to expression systems and methods for expression of desired genes and gene products in cells.

Particularly, the invention relates to a gene encoding a RNA binding protein useful for regulating gene expression in cells, the protein binding site, a gene encoding a regulating protein disulfide isomerase and methods and systems for gene expression of recombinant molecules.

15 Background

Expression systems for expression of exogenous foreign genes in eukaryotic and prokaryotic cells are basic components of recombinant DNA technology. Despite the abundance of expression systems and their wide-spread use, they all have characteristic disadvantages. For example, while expression in *E. coli* is probably the most popular as it is easy to grow and is well understood, eukaryotic proteins expressed therein are not properly modified. Moreover, those proteins tend to precipitate into insoluble aggregates and are difficult to obtain in large amounts. Mammalian expression systems, while practical on small-scale protein production, are more difficult, time-consuming and expensive than in *E. coli*.

A number of plant expression systems exist as well as summarized in US Patent No. 5,234,834, the disclosures of which are hereby incorporated by reference. One advantage of plants or algae in an expression system is that they can be used to produce pharmacologically important proteins and enzymes on a large scale and in relatively pure form. In addition, micro-algae have several unique characteristics that make them ideal organisms for the production of proteins on a large scale.

First, unlike most systems presently used to produce transgenic proteins, algae can be grown in minimal media (inorganic salts) using sunlight as the energy source. These algae can be grown in contained fermentation vessels or on large scale in monitored ponds. Ponds of up to several acres are routinely used for the production of micro-algae. Second, plants and algae have two distinct compartments, the cytoplasm and the chloroplast, in which proteins can be expressed. The cytoplasm of algae is similar to that of other eukaryotic organisms used for protein expression, like yeast and insect cell cultures. The chloroplast is unique to plants and algae and proteins expressed in this environment are likely to have properties different from those of cytoplasmically expressed proteins.

The present invention describes an expression system in which exogenous molecules are readily expressed in either prokaryotic or eukaryotic hosts and in either the cytoplasm or chloroplast. These beneficial attributes are based on the discovery and cloning of components of translation regulation in plants as described in the present invention.

Protein translation plays a key role in the regulation of gene expression across the spectrum of organisms (Kozak, Ann. Rev. Cell Biol., 8:197-225 (1992) and de Smit and Van Duin, Prog. Nucleic Acid Res. Mol. Biol., 38:1-35 (1990)). The majority of regulatory schemes characterized to date involve translational repression often involving proteins binding to mRNA to limit ribosome association (Winter et al., Proc. Natl. Acad. Sci., USA, 84:7822-7826 (1987) and Tang and Draper, Biochem., 29:4434-4439 (1990)). Translational activation has also been observed (Wulczyn and Kahmann, Cell, 65:259-269 (1991)), but few of the underlying molecular mechanisms for this type of regulation have been identified. In plants, light

activates the expression of many genes. Light has been shown to activate expression of specific chloroplast encoded mRNAs by increasing translation initiation (Mayfield et al., Ann. Rev. Plant Physiol. Plant Mol. Biol., 46:147-166 (1995) and Yohn et al., Mol. Cell Biol., 16:3560-3566 (1996)). Genetic evidence in higher plants and algae has shown that nuclear encoded factors are required for translational activation of specific chloroplast encoded mRNAs (Rochaix et al., Embo J., 8:1013-1021 (1989), Kuchka et al., Cell, 58:869-876 (1989), Girard-Bascou et al., Embo J., 13:3170-3181 (1994), Kim et al., Plant Mol. Biol., 127:1537-1545 (1994)).

In the green algae *Chlamydomonas reinhardtii*, a number of nuclear mutants have been identified that affect translation of single specific mRNAs in the chloroplast, often acting at translation initiation (Yohn et al., supra, (1996)). Mutational analysis of chloroplast mRNAs has identified sequence elements within the 5' untranslated region (UTR) of mRNAs that are required for translational activation (Mayfield et al., supra, (1995), Mayfield et al., J. Cell Biol., 127:1537-1545 (1994) and Rochaix, Ann. Rev. Cell Biol., 8:1-28 (1992)), and the 5' UTR of a chloroplast mRNA can confer a specific translation phenotype on a reporter gene *in vivo* (Zerges and Rochaix, Mol. Cell Biol., 14:5268-5277 (1994) and Staub and Maliga, Embo J., 12:601-606 (1993)).

Putative translational activator proteins were identified by purifying a complex of four proteins that binds with high affinity and specificity to the 5' UTR of the chloroplast encoded *psbA* mRNA [encoding the D1 protein, a major component of Photosystem II (PS II)] (Danon and Mayfield, Embo J., 10:3993-4001 (1991)). Binding of these proteins to the 5' UTR of *psbA* mRNA correlates with translation of this mRNA under a variety of

H

physiological (Danon and Mayfield, id., (1991)) and biochemical conditions (Danon and Mayfield, Science, 266:1717-1719 (1994) and Danon and Mayfield, Embo J., 13:2227-2235 (1994)), and in different genetic backgrounds (Yohn et al., supra, (1996)). The binding of this complex to the *psbA* mRNA can be regulated in vitro in response to both redox potential (Danon and Mayfield, Science, 266:1717-1719 (1994)) and phosphorylation (Danon and Mayfield, Embo J., 13:2227-2235 (1994)), both of which are thought to transduce the light signal to activate translation of *psbA* mRNA. The 47 kDa member of the *psbA* RNA binding complex (RB47) is in close contact with the RNA, and antisera specific to this protein inhibits binding to the *psbA* mRNA in vitro (Danon and Mayfield, supra, (1991)).

Although the translational control of *psbA* mRNA by RB47 has been reported, the protein has not been extensively characterized and the gene encoding RB47 has not been identified, cloned and sequenced. In addition, the regulatory control of the activation of RNA binding activity to the binding site by nuclear-encoded trans-acting factors, such as RB60, have not been fully understood. The present invention now describes the cloning and sequencing of both RB47 and RB60. Based on the translation regulation mechanisms of RB47 and RB60 with the RB47 binding site, the present invention also describes a translation regulated expression system for use in both prokaryotes and eukaryotes.

Brief Description of the Invention

The RB47 gene encoding the RB47 activator protein has now been cloned and sequenced, and the target binding site for RB47 on messenger RNA (mRNA) has now been identified. In addition, a regulatory protein disulfide isomerase, a 60 kilodalton protein

referred to as RB60, has also been cloned, sequenced and characterized. Thus, the present invention is directed to gene expression systems in eukaryotic and prokaryotic cells based on translational regulation by RB47 protein, its binding site and the RB60 regulation of RB47 binding site activation.

More particularly, the present invention describes the use of the RB47 binding site, i.e., a 5' untranslated region (UTR) of the chloroplast *psbA* gene, in the context of an expression system for regulating the expression of genes encoding a desired recombinant molecule. Protein translation is effected by the combination of the RB47 binding site and the RB47 binding protein in the presence of protein translation components. Regulation can be further imposed with the use of the RB60 regulatory protein disulfide isomerase. Therefore, the present invention describes reagents and expression cassettes for controlling gene expression by affecting translation of a coding nucleic acid sequence in a cell expression system.

Thus, in one embodiment, the invention contemplates a RB47 binding site sequence, i.e., a mRNA sequence, typically a mRNA leader sequence, which contains the RB47 binding site. A preferred RB47 binding site is *psbA* mRNA. For use in expressing recombinant molecules, the RB47 binding site is typically inserted 5' to the coding region of the preselected molecule to be expressed. In a preferred embodiment, the RB47 binding site is inserted into the 5' untranslated region along with an upstream *psbA* promoter to drive the expression of a preselected nucleic acid encoding a desired molecule. In alternative embodiments, the RB47 binding site is inserted into the regulatory region downstream of any suitable promoter present in a eukaryotic or prokaryotic expression vector. Preferably, the RB47 binding site is positioned within 100 nucleotides of the

translation initiation site. In a further aspect, 3' to the coding region is a 3' untranslated region (3' UTR) necessary for transcription termination and RNA processing.

Thus, in a preferred embodiment, the invention contemplates an expression cassette or vector that contains a transcription unit constructed for expression of a preselected nucleic acid or gene such that upon transcription, the resulting mRNA contains the RB47 binding site for regulation of the translation of the preselected gene transcript through the binding of the activating RB47 protein. The RB47 protein is provided endogenously in a recipient cell and/or is a recombinant protein expressed in that cell.

Thus, the invention also contemplates a nucleic acid molecule containing the sequence of the RB47 gene. The nucleic acid molecule is preferably in an expression vector capable of expressing the gene in a cell for use in interacting with a RB47 binding site. The invention therefore contemplates an expressed recombinant RB47 protein. In one embodiment, the RB47 binding site and RB47 encoding nucleotide sequences are provided on the same genetic element. In alternative embodiments, the RB47 binding site and RB47 encoding nucleotide sequences are provided separately.

The invention further contemplates a nucleic acid molecule containing the sequence encoding the 69 kilodalton precursor to RB47. In alternative embodiments, the RB47 nucleic acid sequence contains a sequence of nucleotides to encode a histidine tag. Thus, the invention relates to the use of recombinant RB47, precursor RB47, and histidine-modified RB47 for use in enhancing translation of a desired nucleic acid.

The invention further contemplates a nucleic acid molecule containing a nucleotide sequence of a polypeptide which

regulates the binding of RB47 to RB47 binding site. A preferred regulatory molecule is the protein disulfide isomerase RB60. The RB60-encoding nucleic acid molecule is preferably in an expression vector capable of expressing the gene in a cell for use in regulating the interaction of RB47 with a RB47 binding site. Thus, the invention also contemplates an expressed recombinant RB60 protein. In one embodiment, the RB47 binding site, RB47 encoding and RB60 encoding nucleotide sequences are provided on the same genetic element. In alternative
10 embodiments, the expression control nucleotide sequences are provided separately. In a further aspect, the RB60 gene and RB47 binding site sequence are provided on the same construct.

The invention can therefore be a cell culture system, an in vitro expression system or a whole tissue, preferably a plant, in which the transcription unit is present that contains the RB47 binding site and further includes a (1) transcription unit capable of expressing RB47 protein or (2) the endogenous RB47 protein itself for the purpose of enhancing translation of the preselected gene having an RB47 binding site in the mRNA.
15

Preferred cell culture systems are eukaryotic and prokaryotic cells. Particularly preferred cell culture systems include plants and more preferably algae.
20

A further preferred embodiment includes (1) a separate transcription unit capable of expressing a regulatory molecule, preferably RB60 protein, or (2) the endogenous RB60 protein itself for the purpose of regulating translation of the preselected gene having an RB47 binding site in the mRNA. In an alternative preferred embodiment, one transcription unit is capable of expressing both the RB47 and RB60 proteins. In a further aspect, the RB47 binding site sequence and RB60 sequence
25
30 are provided on the same construct.

g

In one aspect of the present invention, plant cells endogenously containing RB47 and RB60 proteins are used for the expression of recombinant molecules, such as proteins or polypeptides, through activation of the RB47 binding in an
5 exogenously supplied expression cassette. Alternatively, stable plant cell lines containing endogenous RB47 and RB60 are first generated in which RB47 and/or RB60 proteins are overexpressed. Overexpression is obtained preferably through the stable transformation of the plant cell with one or more expression
10 cassettes for encoding recombinant RB47 and RB60. In a further embodiment, stable cell lines, such as mammalian or bacterial cell lines, lacking endogenous RB47 and/or RB60 proteins are created that express exogenous RB47 and/or RB60.

Plants for use with the present invention can be a
15 transgenic plant, or a plant in which the genetic elements of the invention have been introduced. Based on the property of controlled translation provided by the combined use of the RB47 protein and the RB47 binding site, translation can be regulated for any gene product, and the system can be introduced into any
20 plant species. Similarly, the invention is useful for any prokaryotic or eukaryotic cell system.

Methods for the preparation of expression vectors is well known in the recombinant DNA arts, and for expression in plants is well known in the transgenic plant arts. These particulars
25 are not essential to the practice of the invention, and therefore will not be considered as limiting.

The invention allows for high level of protein synthesis in plant chloroplasts and in the cytoplasm of both prokaryotic and eukaryotic cells. Because the chloroplast is such a productive
30 plant organ, synthesis in chloroplasts is a preferred site of translation by virtue of the large amounts of protein that can

9

be produced. This aspect provides for great advantages in agricultural production of mass quantities of a preselected protein product.

5 The invention further provides for the ability to screen for agonists or antagonists of the binding of RB47 to the RB47 binding site using the expression systems as described herein. Antagonists of the binding are useful in the prevention of plant propagation.

10 Also contemplated by the present invention is a screening assay for agonists or antagonists of RB60 in a manner analogous to that described above for RB47. Such agonists or antagonists would be useful in general to modify expression of RB60 as a way to regulate cellular processes in a redox manner.

15 Kits containing expression cassettes and expression systems, along with packaging materials comprising a label with instructions for use, as described in the claimed embodiments are also contemplated for use in practicing the methods of this invention.

20 Other uses will be apparent to one skilled in the art in light of the present disclosures.

Brief Description of Drawings

In the figures forming a portion of this disclosure:

25 Figures 1A-1D show the complete protein amino acid residue sequence of RB47 is shown from residues 1-623, together with the corresponding nucleic acid sequence encoding the RB47 sequence, from base 1 to base 2732. The nucleotide coding region is shown from base 197-2065, the precursor form. The mature form is from nucleotide position 197-1402. Also shown is the mRNA leader,
30 bases 1-196, and poly A tail of the mRNA, bases 2066-2732. Both the nucleotide and amino acid sequence are listed in SEQ ID NO

be produced. This aspect provides for great advantages in agricultural production of mass quantities of a preselected protein product.

5 The invention further provides for the ability to screen for agonists or antagonists of the binding of RB47 to the RB47 binding site using the expression systems as described herein. Antagonists of the binding are useful in the prevention of plant propagation.

10 Also contemplated by the present invention is a screening assay for agonists or antagonists of RB60 in a manner analogous to that described above for RB47. Such agonists or antagonists would be useful in general to modify expression of RB60 as a way to regulate cellular processes in a redox manner.

15 Kits containing expression cassettes and expression systems, along with packaging materials comprising a label with instructions for use, as described in the claimed embodiments are also contemplated for use in practicing the methods of this invention.

20 Other uses will be apparent to one skilled in the art in light of the present disclosures.

Brief Description of Drawings

In the figures forming a portion of this disclosure:

25 Figures 1A-1D show the complete protein amino acid residue sequence of RB47 is shown from residues 1-623, together with the corresponding nucleic acid sequence encoding the RB47 sequence, from base 1 to base 2732. The nucleotide coding region is shown from base 197-2065, the precursor form. The mature form is from nucleotide position 197-1402. Also shown is the mRNA leader,
30 bases 1-196, and poly A tail of the mRNA, bases 2066-2732. Both the nucleotide and amino acid sequence are listed in SEQ ID NO

5.

Figures 2A-2B show the complete protein amino acid residue sequence of RB60 is shown from residues 1-488, together with the corresponding nucleic acid sequence from base 1 to base 2413, of which bases 16-1614 encode the RB60 sequence. Both the nucleotide and amino acid sequence are listed in SEQ ID NO 10.

Figures 3A-3C show the complete sequence of the *psbA* mRNA, showing both encoded *psbA* protein amino acid residue sequence (residues 1-352) and the nucleic acid sequence as further described in Example 3 is illustrated. Both the nucleotide and amino acid sequence are listed in SEQ ID NO 13.

Figure 4 is a schematic diagram of an expression cassette containing on one transcription unit from 5' to 3', a promoter region derived from the *psbA* gene for encoding the D1 protein from *C. reinhardtii* further containing a transcription initiation site (TS), the RB47 binding site, a region for insertion of a foreign or heterologous coding region, a RB47 coding region, a RB60 coding region, and the 3' flanking region containing transcription termination site (TS), flanked by an origin of replication and selection marker. Restriction endonuclease sites for facilitating insertion of the independent genetic elements are indicated and further described in Example 4A.

Figures 5A-5B show the nucleotide and amino acid sequence of the RB47 molecule containing a histidine tag, the sequences of which are also listed in SEQ ID NO 14.

Figure 6 is a schematic diagram of an expression cassette containing on one transcription unit from 5' to 3', a promoter region derived from the *psbA* gene for encoding the D1 protein from *C. reinhardtii* further containing a transcription initiation site (TS), the RB47 binding site, a region for

AMENDED SHEET 17

IPEAUS 17 AUG 1998

RB47 is also shown in Figures 1A-1D (SEQ ID NO 5). As described in Section 2 above, the predicted protein sequence from the cloned cDNA contained both the derived peptide sequences of RB47 and is highly homologous to poly(A) binding proteins (PABP) from a variety of eukaryotic organisms.

2. Cloning of RB60

To clone the cDNA encoding the 60 kDa *psbA* mRNA binding protein (RB60), the *psbA*-specific RNA binding proteins were purified from light-grown *C. reinhardtii* cells using heparin-agarose chromatography followed by *psbA* RNA affinity chromatography (RAC). RAC-purified proteins were separated by two-dimensional polyacrylamide gel electrophoresis. The region corresponding to RB60 was isolated from the PVDF membrane. RB60 protein was then digested with trypsin. Unambiguous amino acid sequences were obtained from two peptide tryptic fragments (WFVDGELASDYNGPR (SEQ ID NO 6) and (QLILWTTADDLKADAEIMTVFR (SEQ ID NO 7)) as described above for RB47. The calculated molecular weights of the two tryptic peptides used for further analysis precisely matched with the molecular weights determined by mass spectrometry. The DNA sequence corresponding to one peptide of 22 amino acid residues was amplified by PCR using degenerate oligonucleotides, the forward primer 5'CGCGGATCCGAYGCBGAGATYATGAC3' (SEQ ID NO 8) and the reverse primer 5'CGCGAATTCGTCATRATCTCVGCRTC3' (SEQ ID NO 9), where R can be A or G (the other IUPAC nucleotides have been previously defined above). The amplified sequence was then used to screen a λ -gt10 cDNA library

AMENDED SHEET

13

from *C. reinhardtii*. Three clones were identified with the largest being 2.2 kb. Selection and sequencing was performed as described for RB47 cDNA.

The resulting RB60 cDNA sequence is available via
5 GenBank (Accession Number AF027727). The nucleotide and encoded amino acid sequence of RB60 is also shown in Figures 2A-2B (SEQ ID NO 10). The protein coding sequence of 488 amino acid residues corresponds to nucleotide positions 16-1614 of the 2413 base pair
10 sequence. The predicted amino acid sequence of the cloned cDNA contained the complete amino acid sequences of the two tryptic peptides. The amino acid sequence of the encoded protein revealed that it has high sequence homology to both plant and mammalian protein disulfide
15 isomerase (PDI), and contains the highly conserved thioredoxin-like domains with -CysGlyHisCys- (-CGHC-) (SEQ ID NO 11) catalytic sites in both the N-terminal and C-terminal regions and the -LysAspGluLeu- (-KDEL-) (SEQ ID NO 12) endoplasmic reticulum (ER) retention
20 signal at the C-terminus found in all PDIs. PDI is a multifunctional protein possessing enzymatic activities for the formation, reduction, and isomerization of disulfide bonds during protein folding, and is typically found in the ER. The first 30 amino acid residues of
25 RB60 were found to lack sequence homology with the N-terminal signal sequence of PDI from plants or mammalian cells. However, this region has characteristics of chloroplast transit peptides of *C. reinhardtii*, which have similarities with both
30 mitochondrial and higher plant chloroplast presequences. A transit peptide sequence should override the function

of the -KDEL- ER retention signal and target the protein to the chloroplast since the -KDEL- signal acts only to retain the transported protein in the ER.

5 3. Preparation of psbA Promoter Sequence and RB47
Binding Site Nucleotide Sequence

10 The chloroplast *psbA* gene from the green unicellular alga *C. reinhardtii* was cloned and sequenced as described by Erickson et al., Embo J., 3:2753-2762 (1984), the disclosure of which is hereby incorporated by reference. The DNA sequence of the coding regions and the 5' and 3' untranslated (UTR) flanking sequences of the *C. reinhardtii psbA* gene is shown in Figures 3A-3C. The *psbA* gene sequence is also available through 15 GenBank as further discussed in Example 4. The nucleotide sequence is also listed as SEQ ID NO 13. The deduced amino acid sequence (also listed in SEQ ID NO 13) of the coding region is shown below each codon beginning with the first methionine in the open reading 20 frame. Indicated in the 5' non-coding sequence are a putative Shine-Dalgarno sequence in the dotted box, two putative transcription initiation sites determined by S1 mapping (S1) and the Pribnow-10 sequence in the closed box. Inverted repeats of eight or more base pairs are 25 marked with arrows and labeled A-D. A direct repeat of 31 base pairs with only two mismatches is marked with arrows labeled 31. Indicated in the 3' non-coding sequence is a large inverted repeat marked by a forward arrow and the S1 cleavage site marking the 3' end of the 30 mRNA. Both the 5' and 3' untranslated regions are used in preparing one of the expression cassettes of this

invention as further described below.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
221

The above-identified regions are used to prepare expression constructs as described below. The promoter and RB47 binding site regions can be used separately; for example, the RB47 binding site sequence can be isolated and used in a eukaryotic or prokaryotic plasmid with a non-*psbA* promoter. Alternatively, the entire *psbA* 5' UTR having 251 nucleotides as shown in Figures 3A-3C is used for the regulatory region in an expression cassette containing both the *psbA* promoter and RB47 binding site sequence as described below.

4. Preparation of Expression Vectors and Expression of Coding Sequences

A. Constructs Containing an *psbA* Promoter, an RB47 Binding Site Nucleotide Sequence, a Desired Heterologous Coding Sequence, an RB47-Encoding Sequence and an RB60-Encoding Sequence

Plasmid expression vector constructs, alternatively called plasmids, vectors, constructs and the like, are constructed containing various combinations of elements of the present invention as described in the following examples. Variations of the positioning and operably linking of the genetic elements described in the present invention and in the examples below are contemplated for use in practicing the methods of this invention. Methods for manipulating DNA elements into operable expression cassettes are well known in the art of molecular biology. Accordingly, variations of control elements, such as constitutive or inducible promoters, with respect to prokaryotic or

eukaryotic expression systems as described in Section C. are contemplated herein although not enumerated.

Moreover, the expression the various elements is not limited to one transcript producing one mRNA; the

5 invention contemplates protein expression from more than one transcript if desired.

As such, while the examples below recite one or two types of expression cassettes, the genetic elements of RB47 binding site, any desired coding sequence, in
10 combination with RB47 and RB60 coding sequences along with a promoter are readily combined in a number of operably linked permeations depending on the requirements of the cell system selected for the expression. For example, for expression in a
15 chloroplast, endogenous RB47 protein is present therefore an expression cassette having an RB47 binding site and a desired coding sequence is minimally required along with an operative promoter sequence.

Overexpression of RB47 may be preferable to enhance the
20 translation of the coding sequence; in that case, the chloroplast is further transformed with an expression cassette containing an RB47-encoding sequence. Although the examples herein and below utilize primarily the sequence encoding the precursor form of RB47, any of the
25 RB47-encoding sequences described in the present invention, i.e., RB47 precursor, mature RB47 and histidine-modified RB47 are contemplated for use in any expression cassette and system as described herein. To regulate the activation of translation, an RB60-encoding
30 element is provided to the expression system to provide the ability to regulate redox potential in the cell as

AMENDED SHEET
18

taught in Section B. These examples herein and below represent a few of the possible permutations of genetic elements for expression in the methods of this invention.

5 In one embodiment, a plasmid is constructed containing an RB47 binding site directly upstream of an inserted coding region for a heterologous protein of interest, and the RB47 and RB60 coding regions. Heterologous refers to the nature of the coding region
10 being dissimilar and not from the same gene as the regulatory molecules in the plasmid, such as RB47 and RB60. Thus, all the genetic elements of the present invention are produced in one transcript from the IPTG-inducible *psbA* promoter. Alternative promoters are
15 similarly acceptable.

The final construct described herein for use in a prokaryotic expression system makes a single mRNA from which all three proteins are translated. The starting plasmid is any *E. coli* based plasmid containing an
20 origin of replication and selectable marker gene. For this example, the Bluescript plasmid, pBS, commercially available through Stratagene, Inc., La Jolla, CA, which contains a polylinker-cloning site and an ampicillin resistant marker is selected for the vector.

25 The wild-type or native *psbA* gene (Erickson et al., Embo J., 3:2753-2762 (1984), also shown in Figures 3A-3C, is cloned into pBS at the EcoRI and BamHI sites of the polylinker. The nucleotide sequence of the *psbA* gene is available on GenBank with the 5' UTR and 3' UTR
30 respectively listed in Accession Numbers X01424 and X02350. The EcoRI site of *psbA* is 1.5 kb upstream of

the *psbA* initiation codon and the BamHI site is 2 kb downstream of the stop codon. This plasmid is referred to as pDl.

Using site-directed PCR mutagenesis, well known to one of ordinary skill in the art, an NdeI site is placed at the initiation codon of *psbA* in the pDl plasmid so that the ATG of the NdeI restriction site is the ATG initiation codon. This plasmid is referred to as pDl/Nde. An Nde site is then placed at the initiation codon of the gene encoding the heterologous protein of interest and an Xho I site is placed directly downstream (within 10 nucleotides) of the TAA stop codon of the heterologous protein coding sequence. Again using site-directed mutagenesis, an XhoI site is placed within 10 nucleotides of the initiation codon of RB47, the preparation of which is described in Example 2, and an NotI site is placed directly downstream of the stop codon of RB47. The heterologous coding region and the RB47 gene are then ligated into pDl/Nde so that the heterologous protein gene is directly adjacent to the RB47 binding site and the RB47 coding region is downstream of the heterologous coding region, using the Xho I site at the heterologous stop codon and the Not I site of the pDl polylinker.

These genetic manipulations result in a plasmid containing the 5' end of the *psbA* gene including the promoter region and with the RB47 binding site immediately upstream of a heterologous coding region, and the RB47 coding region immediately downstream of the heterologous coding region. The nucleotides between the stop codon of the heterologous coding region and the

initiation codon of the RB47 coding region is preferably less than 20 nucleotides and preferably does not contain any additional stop codons in any reading frame. This plasmid is referred to as pD1/RB47.

5 Using site-directed mutagenesis, a NotI site is placed immediately (within 10 nucleotides) upstream of the initiation codon of RB60, the preparation of which is described in Example 2, and an Xba I site is placed downstream of the RB60 stop codon. This DNA fragment is then ligated to the 3' end of the *psbA* gene using the Xba I site found in the 3' end of the *psbA* gene so that the *psbA* 3' end is downstream of the RB60 coding region. This fragment is then ligated into the pD1/RB47 plasmid using the NotI and BamHI sites so that the RB60 coding region directly follows the RB47 coding region. The resulting plasmid is designated pD1/RB47/RB60. Preferably there is less than 20 nucleotides between the RB47 and RB60 coding regions and preferably there are no stop codons in any reading frame in that region. The final plasmid thus contains the following genetic elements operably linked in the 5' to 3' direction: the 5' end of the *psbA* gene with a promoter capable of directing transcription in chloroplasts, an RB47 binding site, a desired heterologous coding region, the RB47 coding region, the RB60 coding region, and the 3' end of the *psbA* gene which contains a transcription termination and mRNA processing site, and an *E. coli* origin of replication and ampicillin resistance gene. A diagram of this plasmid with the restriction sites is shown in Figure 4.

Expression of pD1/RB47/RB60 in *E. coli* to produce

recombinant RB47, RB60 and the recombinant heterologous protein is performed as described in Example 4B. The heterologous protein is then purified as further described.

5 Expression cassettes in which the sequences encoding RB47 and RB60 are similarly operably linked to a heterologous coding sequence having the *psbA* RB47 binding site as described in Example 3 are prepared with a different promoter for use in eukaryotic, such as
10 mammalian expression systems. In this aspect, the cassette is similarly prepared as described above with the exception that restriction cloning sites are dependent upon the available multiple cloning sites in the recipient vector. Thus, the RB47 binding site
15 prepared in Example 3 is prepared for directed ligation into a selected expression vector downstream of the promoter in that vector. The RB47 and RB60 coding sequences are obtained from the pD1/RB47/RB60 plasmid by digestion with XhoI and XbaI and inserted into a
20 similarly digested vector if the sites are present. Alternatively, site-directed mutagenesis is utilized to create appropriate linkers. A desired heterologous coding sequence is similarly ligated into the vector for expression.

25

B. Constructs Containing RB47 Nucleotide Sequence

1) Purified Recombinant RB47 Protein

In one approach to obtain purified recombinant RB47 protein, the full length RB47 cDNA
30 prepared above was cloned into the *E. coli* expression vector pET3A (Studier et al., Methods Enzymol., 185:60-

89 (1990)), also commercially available by Novagen, Inc., Madison, WI and transformed into BL21 *E. coli* cells. The cells were grown to a density of 0.4 (OD_{600}), then induced with 0.5 mM IPTG. Cells were then allowed to grow for an additional 4 hours, at which point they were pelleted and frozen.

Confirmation of the identity of the cloned cDNA as encoding the authentic RB47 protein was accomplished by examining protein expressed from the cDNA by immunoblot analysis and by RNA binding activity assay. The recombinant RB47 protein produced when the RB47 cDNA was expressed was recognized by antisera raised against the *C. reinhardtii* RB47 protein. The *E. coli* expressed protein migrated at 80 kDa on SDS-PAGE, but the protein was actually 69 kDa, as determined by mass spectrometry of the *E. coli* expressed protein. This mass agrees with the mass predicted from the cDNA sequence. A 60 kDa product was also produced in *E. coli*, and recognized by the antisera against the *C. reinhardtii* protein, which is most likely a degradation or early termination product of the RB47 cDNA. The recombinant RB47 protein expressed from the RB47 cDNA is recognized by the antisera raised against the *C. reinhardtii* protein at levels similar to the recognition of the authentic *C. reinhardtii* RB47 protein, demonstrating that the cloned cDNA produces a protein product that is immunologically related to the naturally produced RB47 protein. In order to generate a recombinant equivalent of the endogenous native RB47, the location of the 47 kDa polypeptide was mapped on the full-length recombinant protein by comparing mass spectrometric data of tryptic

IPEAUS 17 AUG 1998

digests of the *C. reinhardtii* 47 kDa protein and the full-length recombinant protein. Thus, peptide mapping by mass spectrometry has shown that the endogenous RB47 protein corresponds primarily to the RNA binding domains contained within the N-terminal region of the predicted precursor protein, suggesting that a cleavage event is necessary to produce the mature 47 kDa protein. Thus, full-length recombinant RB47 is 69 kDa and contains a carboxy domain that is cleaved *in vivo* to generate the endogenous mature form of RB47 that is 47 kDa.

To determine if the heterologously expressed RB47 protein was capable of binding the *psbA* RNA, the *E. coli* expressed protein was purified by heparin agarose chromatography. The recombinant RB47 protein expressed in *E. coli* was purified using a protocol similar to that used previously for purification of RB47 from *C. reinhardtii*. Approximately 5 g of *E. coli* cells grown as described above were resuspended in low salt extraction buffer (10 mM Tris [pH 7.5], 10 mM NaCl, 10 mM MgCl₂, 5 mM β -mercaptoethanol) and disrupted by sonication. The soluble cell extract was applied to a 5 mL Econo-Pac heparin cartridge (Bio-Rad) which was washed prior to elution of the RB47 protein (Danon and Mayfield, Embo J., 10:3993-4001 (1991)).

The *E. coli* expressed protein that bound to the heparin agarose matrix was eluted from the column at the same salt concentration as used to elute the authentic *C. reinhardtii* RB47 protein. This protein fraction was used in *in vitro* binding assays with the *psbA* 5' UTR.

Both the 69 and 60 kDa *E. coli* expressed proteins crosslinked to the radiolabeled *psbA* 5' UTR at levels

PEAUS 17 AUG 1998

similar to crosslinking of the endogenous RB47 protein, when the RNA/protein complex is subjected to UV irradiation.

Heparin agarose purified proteins, both from the *E. coli* expressed RB47 cDNA and from *C. reinhardtii* cells, were used in an RNA gel mobility shift assay to determine the relative affinity and specificity of these proteins for the 5' UTR of the *psbA* mRNA. The *E. coli* expressed proteins bound to the *psbA* 5' UTR *in vitro* with properties that are similar to those of the endogenous RB47 protein purified from *C. reinhardtii*. RNA binding to both the *E. coli* expressed and the endogenous RB47 protein was competed using either 200 fold excess of unlabeled *psbA* RNA or 200 fold excess of poly(A) RNA. RNA binding to either of these proteins was poorly competed using 200 fold excess of total RNA or 200 fold excess of the 5' UTR of the *psbD* or *psbC* RNAs. Different forms of the RB47 protein (47 kDa endogenous protein vs. the 69 kDa *E. coli* expressed protein) may account for the slight differences in mobility observed when comparing the binding profiles of purified *C. reinhardtii* protein to heterologously expressed RB47.

The mature form of RB47 is also produced in recombinant form by the insertion by PCR of an artificial stop codon in the RB47 cDNA at nucleotide positions 1403-1405 with a stop codon resulting in a mature RB47 recombinant protein having 402 amino acids as shown in Figures 1A-1D. An example of this is shown in Figures 5A-5B for the production of a recombinant histidine-modified RB47 mature protein as described

below. The complete RB47 cDNA is inserted into an expression vector, such as pET3A as described above, for expression of the mature 47 kDa form of the RB47 protein. In the absence of the inserted stop codon, the transcript reads through to nucleotide position 2066-2068 at the TAA stop codon to produce the precursor RB47 having the above-described molecular weight characteristics and 623 amino acid residues.

Recombinant RB47 is also expressed and purified in plant cells. For this aspect, *C. reinhardtii* strains were grown in complete media (Tris-acetate-phosphate [TAP] (Harris, The *Chlamydonas* Sourcebook, San Diego, CA, Academic Press (1989)) to a density of 5×10^6 cells/mL under constant light. Cells were harvested by centrifugation at 4°C for 5 minutes at 4,000 g. Cells were either used immediately or frozen in liquid N₂ for storage at -70°C.

Recombinant RB47 protein was also produced as a modified RB47 protein with a histidine tag at the amino-terminus according to well known expression methods using pET19-D vectors available from Novagen, Inc., Madison, WI. The nucleotide and amino acid sequence of a recombinant histidine-modified RB47 of the mature 47 kDa form is shown in Figures 5A-5B with the nucleotide and amino acid sequence also listed in SEQ ID NO 14. Thus the nucleotide sequence of a histidine-modified RB47 is 1269 bases in length. The precursor form of the RB47 protein is similarly obtained in the expression system, both of which are modified by the presence of a histidine tag that allows for purification by metal affinity chromatography.

IPEAUS 17 AUG 1998

The recombinant histidine-modified RB47 purified through addition of a poly-histidine tag followed by Ni^{+2} column chromatography showed similar binding characteristics as that described for recombinant precursor RB47 described above.

C. Constructs Containing RB60 Nucleotide Sequence

In one approach to obtain purified recombinant RB60 protein, the full-length RB60 cDNA prepared above was cloned into the *E. coli* expression vector pET3A (Studier et al., Methods Enzymol., 185:60-89 (1990)), also commercially available by Novagen, Inc., Madison, WI and transformed into BL21 *E. coli* cells. The cells were grown to a density of 0.4 (OD_{600}), then induced with 0.5 mM IPTG. Cells were then allowed to grow for an additional 4 hours, at which point they were pelleted and frozen.

Recombinant histidine-modified RB60 was also expressed with a pET19-D vector as described above for RB47 that was similarly modified. Purification of the recombinant RB60 proteins was performed as described for RB47 thereby producing recombinant RB60 proteins for use in the present invention.

The RB60 coding sequence is also mutagenized for directional ligation into an selected vector for expression in alternative systems, such as mammalian expression systems.

D. Constructs Containing an RB47-Encoding Sequence and an RB60-Encoding Sequence

To prepare an expression cassette for encoding

IPEAUS 17 AUG 1998

both RB47 and RB60, one approach is to digest plasmid pD1/RB47/RB60 prepared above with XhoI and XbaI to isolate the fragment for both encoding sequences. The fragment is then inserted into a similarly digested expression vector if available or is further mutagenized to prepare appropriate restriction sites.

Alternatively, the nucleotide sequences of RB47 and RB60, as described in Example 2, are separately prepared for directional ligation into a selected vector.

An additional embodiment of the present invention is to prepare an expression cassette containing the RB47 binding site along with the coding sequences for RB47 and RB60, the plasmid pD1/RB47/RB60 prepared above is digested with NdeI and XhoI to prepare an expression cassette in which any desired coding sequence having similarly restriction sites is directionally ligated. Expression vectors containing both the RB47 and RB60 encoding sequences in which the RB47 binding site sequence is utilized with a different promoter are also prepared as described in Example 4A.

E. Constructs Containing an RB47 Binding Site Nucleotide Sequence, Insertion Sites for a Desired Heterologous Coding Sequence, and an RB47-Encoding Sequence

In another permutation, a plasmid or expression cassette is constructed containing an RB47 binding site directly upstream of an inserted coding region for a heterologous protein of interest, and the RB47 coding region. The final construct described herein for use in a prokaryotic expression system makes

IPEAUS 17 AUG 1998

a single mRNA from which both proteins are translated.

The plasmid referred to as pD1/RB47 is prepared as described above in Example 4A. A diagram of this plasmid with the restriction sites is shown in Figure 6.

5 Expression of pD1/RB47 in *E. coli* to produce recombinant RB47 and the recombinant heterologous protein is performed as described in above. The heterologous protein is then purified as further described.

10 To produce an expression cassette that allows for insertion of an alternative desired coding sequence, the plasmid pD1/RB47 is digested with NdeI and XhoI resulting in a vector having restriction endonuclease sites for insertion of a desired coding sequence
15 operably linked to a RB47 binding site and RB47 coding sequence on one transcriptional unit.

20 F. Constructs Containing an RB47 Binding Site Nucleotide Sequence, Insertion Sites for a Desired Heterologous Coding Sequence, and an RB47-Encoding Sequence

In another permutation, a plasmid or expression cassette is constructed containing an RB47 binding site directly upstream of an inserted coding
25 region for a heterologous protein of interest, and the RB60 coding region. The final construct described herein for use in a prokaryotic expression system makes a single mRNA from which both proteins are translated. In this embodiment, a separate construct encoding
30 recombinant RB47 as described in Example 4B is co-transformed into the *E. coli* host cell for expression.

The plasmid referred to as pD1/RB60 is prepared as described above for pD1/RB47 in Example 4A with the exception that XhoI and XbaI sites are created on RB60 rather than RB47.

5 Expression of pD1/RB60 in *E. coli* to produce recombinant RB60 and the recombinant heterologous protein is performed as described in above with the combined expression of RB47 from a separate expression cassette. The heterologous protein is then purified as
10 further described.

To produce an expression cassette that allows for insertion of an alternative desired coding sequence, the plasmid pD1/RB60 is digested with NdeI and XhoI
15 resulting in a vector having restriction endonuclease sites for insertion of a desired coding sequence operably linked to a RB47 binding site and RB60 coding sequence on one transcriptional unit.

20 G. Constructs Containing RB47 Binding Site
Nucleotide Sequence and Heterologous Coding
Sequences

1) Expression of Recombinant Tetanus Toxin
Single Chain Antibody

The examples herein describe constructs
25 that are variations of those described above. The constructs described below contain an RB47 binding site sequence and a heterologous coding sequence. The activating protein RB47 was endogenously provided in the chloroplast and or plant cell. In other aspects however
30 as taught by the methods of the present invention, the chloroplast is further transformed with an RB47-

expression construct as described above for overexpression of RB47 to enhance translation capacities.

A strain of the green algae *Chlamydomonas reinhardtii* was designed to allow expression of a single chain antibody gene in the chloroplast. The transgenically expressed antibody was produced from a chimeric gene containing the promoter and 5' untranslated region (UTR) of the chloroplast *psbA* gene prepared as described above, followed by the coding region of a single chain antibody (encoding a tetanus toxin binding antibody), and then the 3' UTR of the *psbA* gene also prepared as described above to provide for transcription termination and RNA processing signals. This construct is essentially pD1/Nde including a heterologous coding sequence having a 3' XbaI restriction site for ligation with the 3' *psbA* gene and is diagramed in Figure 7.

The *psbA*-single chain construct was first transformed into *C. reinhardtii* chloroplast and transformants were then screened for single chain gene integration. Transformation of chloroplast was performed via bolistic delivery as described in US Patents 5,545,818 and 5,553,878, the disclosures of which are hereby incorporated by reference. Transformation is accomplished by homologous recombination via the 5' and 3' UTR of the *psbA* mRNA.

As shown in Figure 8, two of the transformants that contained the single chain chimeric gene produced single chain antibodies at approximately 1% of total protein levels. The transgenic antibodies were of the correct

size and were completely soluble, as would be expected of a correctly folded protein. Few degradation products were detectable by this Western analysis, suggesting that the proteins were fairly stable within the chloroplast. To identify if the produced antibody retained the binding capacity for tetanus toxin, ELISA assays were performed using a mouse-produced Fab, from the original tetanus toxin antibody, as the control. The chloroplast single chain antibody bound tetanus toxin at levels similar to Fab, indicating that the single chain antibody produced in *C. reinhardtii* is a fully functional antibody. These results clearly demonstrate the ability of the chloroplast to synthesis and accumulate function antibody molecules resulting from the translational activation of an RB47 binding site in an expression cassette by endogenous RB47 protein in the chloroplast.

2) Expression of Bacterial Luciferase Enzyme Having Two Subunits

For the production of molecules that contain more than one subunit, such as dIgA and bacterial luciferase enzyme, several proteins must be produced in stoichiometric quantities within the chloroplast. Chloroplast have an advantage for this type of production over cytoplasmic protein synthesis in that translation of multiple proteins can originate from a single mRNA. For example, a dicistronic mRNA having 5' and 3' NdeI and XbaI restriction sites and containing both the A and B chains of the bacterial luciferase enzyme was inserted downstream of the *psbA* promoter and

5' UTR of the pD1/Nde construct prepared in Example 4A above. In this construct, the bacterial LuxAB coding region was ligated between the *psbA* 5' UTR and the *psbA* 3' end in an *E. coli* plasmid that was then transformed into *Chlamydomonas reinhardtii* cells as described above for expression in the chloroplast. A schematic of the construct is shown in Figure 9. Single transformant colonies were then isolated. A plate containing a single isolate was grown for 10 days on complete media and a drop of the luciferase substrate n-Decyl Aldehyde was placed on the plate and the luciferase visualized by video-photography in a dark chamber. Both proteins were synthesized from this single mRNA and luciferase activity accumulated within the chloroplast as shown in Figure 10. Some mRNA within plastids contained as many as 5 separate proteins encoded on a single mRNA.

3) Expression of Dimeric IgA

To generate dimeric IgA, the construct shown in Figure 11 is engineered so that the *psbA* promoter and 5' UTR are used to drive the synthesis of the light chain and heavy chains of an antibody, and the J chain normally associated with IgA molecules. The nucleic acid sequences for the dimeric IgA are inserted into the RB47 binding site construct prepared in Example 4A. The construct is then transformed into *C. reinhardtii* cells as previously described for expression of the recombinant dIgA.

Production of these three proteins within the plastid allows for the self assembly of a dimeric IgA (dIgA). Production of this complex is monitored in

several ways. First, Southern analysis of transgenic algae is used to identify strains containing the polycistronic chimeric dIgA gene. Strains positive for integration of the dIgA gene are screened by Northern
5 analysis to ensure that the chimeric mRNA is accumulating. Western blot analysis using denaturing gels is used to monitor the accumulation of the individual light, heavy and J chain proteins, and native
10 gels Western blot analysis will be used to monitor the accumulation of the assembled dIgA molecule.

By using a single polycistronic mRNA in the context of RB47 regulated translation, two of the potential pitfalls in the assembly of multimeric dIgA molecule are overcome. First, this construct ensures approximately
15 stoichiometric synthesis of the subunits, as ribosomes reading through the first protein are likely to continue to read through the second and third proteins as well. Second, all of the subunits are synthesized in close physical proximity to each other, which increases the
20 probability of the proteins self assembling into a multimeric molecule. Following the production of a strain producing dIgA molecules, the production of dIgA on an intermediate scale by growing algae in 300 liter fermentors is then performed. Larger production scales
25 are then performed thereafter.

The foregoing specification, including the specific embodiments and examples, is intended to be illustrative of the present invention and is not to be taken as
30 limiting. Numerous other variations and modifications can be effected without departing from the true spirit and scope of the invention.

A11 B1

AMENDED SHEET 34